

PATENT COOPERATION TREATY

PCT

NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

United States Patent and Trademark
Office
(Box PCT)
Crystal Plaza 2
Washington, DC 20231
ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

Date of mailing (day/month/year) 20 July 1999 (20.07.99)	
International application No. PCT/GB98/03687	Applicant's or agent's file reference PPD 50288/WO
International filing date (day/month/year) 10 December 1998 (10.12.98)	Priority date (day/month/year) 11 December 1997 (11.12.97)
Applicant JEPSON, Ian et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:
25 May 1999 (25.05.99)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was

☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland Facsimile No.: (41-22) 740.14.35	Authorized officer <p style="text-align: center;">C. Carrié</p> Telephone No.: (41-22) 338.83.38
--	--

REC'D 18 FEB 2000

WIPO PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference PPD 50288/WO	FOR FURTHER ACTION See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416)	
International application No. PCT/GB98/03687	International filing date (day/month/year) 10/12/1998	Priority date (day/month/year) 11/12/1997
International Patent Classification (IPC) or national classification and IPC C12N15/82		
Applicant ZENECA LIMITED et al.		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.



2. This REPORT consists of a total of 5 sheets, including this cover sheet.

- ☐ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☐ Priority
- III ☒ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand 25/05/1999	Date of completion of this report 14.02.00
Name and mailing address of the international preliminary examining authority:  European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465	Authorized officer Keller, Y Telephone No. +49 89 2399 7419 

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**

International application No. PCT/GB98/03687

I. Basis of the report

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

Description, pages:

1-19 as originally filed

Claims, No.:

1-17 as originally filed

Drawings, sheets:

1/16-16/16 as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description, pages:
☐ the claims, Nos.:
☐ the drawings, sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

III. Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

The questions whether the claimed invention appears to be novel, to involve an inventive step (to be non-obvious), or to be industrially applicable have not been examined in respect of:

- ☐ the entire international application.
☒ claims Nos. 17.

because:

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT**International application No. **PCT/GB98/08687**

- ☐ the said international application, or the said claims Nos. relate to the following subject matter which does not require an international preliminary examination (*specify*):
- ☒ the description, claims or drawings (*indicate particular elements below*) or said claims Nos. 17 are so unclear that no meaningful opinion could be formed (*specify*):
- see separate sheet**
- ☐ the claims, or said claims Nos. are so inadequately supported by the description that no meaningful opinion could be formed.
- ☐ no international search report has been established for the said claims Nos. .

V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**1. Statement**

Novelty (N)	Yes:	Claims 9, 10, 13, 14
	No:	Claims 1-8; 11, 12, 15, 16
Inventive step (IS)	Yes:	Claims
	No:	Claims 1-16
Industrial applicability (IA)	Yes:	Claims 1-16
	No:	Claims

2. Citations and explanations**see separate sheet****VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

see separate sheet

Re Item III

Non-establishment of opinion with regard to novelty, inventive step and industrial applicability

Claim 17 is a so called "omnibus claim" which is devoid of any technical feature necessary to properly define the claimed subject-matter.

Therefore, a meaningful evaluation with regard to novelty, inventive step and industrial applicability can not be carried out.

Re Item V

Reasoned statement under Rule 66.2(a)(ii) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement

Reference is made to the following documents:

D1: EP 0442 592

D2: WO 9707221

D3: WO 9321334

Novelty :

1. D1 discloses a DNA sequence containing genes involved in the sucrose metabolism or distribution (column 3, lines 1-11) such as e.g. invertase (Example 3, claims). These genes are placed under control of promoters, such promoters can be constitutive, or tissue specific. The transformation of plants with such constructs is used to increase the yield of such plants (column 1, lines 39-48; column 18, lines 15-17; Example 3).
2. D2 discloses also the use of DNA constructs in a method to increase the yield of plants by transforming plants or cells with said DNA constructs (Abstract). Such DNA constructs contain gene(s) involved in the sucrose metabolism such as invertase and hexokinase (page 1, § 1). Said genes are under the control of

**INTERNATIONAL PRELIMINARY
EXAMINATION REPORT - SEPARATE SHEET**

International application No. PCT/GB98/03687

promoter(s). These promoters can be tissue specific promoters, promoters activated during certain development stages or promoters which can be induced time and spacially induced by external agents such as chemicals (page 8 last § to page 10, § 1; claims 12, 13).

3. D1 and D2 disclose methods to increase yields of plants or flowering, by transforming plants with genes involved in the sucrose metabolism said genes being linked to controllable promoters. The teaching of D1 and/or D2 does not differs from the subject-matter of claims 1-8, 11, 12, 15, 16 especially in regard to the claimed promoter for use in these methods since the claims are drafted in a rather broad way (see Item VIII).

Thus, in view of D1 and/or D2 claims 1-8, 11, 12, 15, 16 do not meet the requirements of Article 33(2) PCT.

Inventive step :

4. The subject-matter of claims 9-10, 13-14 merely consist in the selection of a particular chemically inducible promoter system (disclosed in D3, page 8 line 20 to page 9, line 2). Such a selection can only be regarded as involving an inventive step if a surprising or unexpected effect, for the skilled person, can be shown. This is not the case for the present claimed subject-matter.

Therefore, said claims do not meet the requirement of Article 33(3) PCT.

Re Item VIII

Certain observations on the international application

1. Claims 1-5 do not contain the essential technical feature for the definition of the subject-matter. The claims attempt to define the subject-matter in terms of the result to be achieved which merely amounts to a statement of the underlying problem. The technical features necessary for achieving this result should be added.

Hence, claims 1-5 do not meet the requirements of Article 6 PCT.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, 9/26, A01H 5/00	A1	(11) International Publication Number: WO 99/29881 (43) International Publication Date: 17 June 1999 (17.06.99)
(21) International Application Number: PCT/GB98/03687 (22) International Filing Date: 10 December 1998 (10.12.98) (30) Priority Data: 97121829.2 11 December 1997 (11.12.97) EP (71) Applicant (for all designated States except US): ZENECA LIMITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): JEPSON, Ian [GB/GB]; Jealott's Hill Research Station, Bracknell, Berkshire RG42 6ET (GB). CHU, Chengcai [CN/DE]; Institute of Plant Genetics and Crop, Plant Research (IPK), Corrensstrasse 3, D-06466 Gatersleben (DE). QU, Nan [CN/DE]; Institute of Plant Genetics and Crop, Plant Research (IPK), Corrensstrasse 3, D-06466 Gatersleben (DE). SONNEWALD, Uwe [DE/DE]; Institute of Plant Genetics and Crop, Plant Research (IPK), Corrensstrasse 3, D-06466 Gatersleben (DE).		(74) Agents: HOUGHTON, Malcolm, John et al.; Zeneca Agrochemicals, Intellectual Property Dept., Jealotts Hill Research Station, P.O. Box 3538, Bracknell, Berkshire RG42 6YA (GB). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>

(54) Title: GENETIC METHOD**(57) Abstract**

The present invention describes a method of increasing plan yield. Also described are DNA constructs comprising DNA sequences coding for proteins involved in sucrose transport, metabolism and uptake operably linked to controllable promoter regions and plants transformed with said constructs. More particularly a method for the controlled production of said proteins resulting in an alteration in plant growth characteristics, flowering time and in yield is described.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

GENETIC METHOD

The present invention relates to a method of increasing plant yield, to DNA constructs comprising DNA sequences coding for proteins involved in sucrose transport, metabolism and uptake operably linked to controllable promoter regions and to plants transformed with said constructs. More particularly the present invention relates to the controlled production of said proteins resulting in an alteration in plant growth characteristics, flowering time and in yield.

Photosynthesis is the major source of energy used to support biological processes in higher plants. The photosynthesising cells serve as important sources of photoassimilates or organic compounds produced in the plant by photosynthesis. Most fixed organic carbon is translocated from the source photosynthetic tissue to the non-photosynthetic organs which are known as the sink and this is the area in the plant where the translocated nutrients are either used or stored. The principal product of carbon fixation during the photosynthetic reaction is the disaccharide sucrose.

We have now found that by controlling the expression of DNA sequences coding for proteins involved in the transport, metabolism and uptake of sucrose using inducible promoter systems, it is possible to alter the sucrose levels in the plant in a controlled manner to produce the desired change in flowering and/or plant weight and/or height at the appropriate stage in plant growth whereby any effects deleterious to the plant are avoided and the overall yield of the plant is increased. The use of controllable promoter regions permits the expression of said DNA sequences to be regulated in a very precise way such that the optimal level of expression, the optimal time at which the DNA sequence is expressed and the optimal location in the plant may be determined.

According to a first aspect of the present invention there is provided a method of increasing the yield of a plant comprising transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to a controllable promoter region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence(s) from said controllable promoter

region by application of an external chemical inducer whereby the yield of said transgenic plant is increased.

According to a preferred embodiment of the first aspect of the present invention there is provided a method of increasing the yield of a plant by selectively increasing the importation of fixed carbon into photosynthetically inactive sink tissues comprising
5 transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to a controllable promoter region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA
10 sequence(s) from said controllable promoter region by application of an external chemical inducer whereby the transportation of fixed carbon from photosynthetically active source tissue to photosynthetically inactive sink tissue of said transgenic plant is selectively increased.

As used herein the term "source tissue" is used to denote those photosynthetically
15 active tissues of the plant which are net exporters of fixed carbon and "sink tissue" denotes those photosynthetically inactive tissues of the plant which are net importers of fixed carbon.

It is economically and practically very desirable to be able to control both the ability to flower and the time of flowering of a plant. In some instances it may be desirable to synchronise flowering or to switch on flowering early or to manipulate flowering behaviour
20 to suit the constraints imposed by growing in particular geographical areas. Generally an increase in the number of flowers is reflected in an increase in the eventual yield from the plant due to the increase in the number of seeds.

Similarly, an increase in the fresh weight of a plant as measured by an increase in leaf area results in an increase in yield due to the increase in the photosynthetic capacity of the
25 plant.

Yield depends upon at least two parameters:- (i) sink induction and (ii) sink growth. Amongst other factors, sink induction can be stimulated by reducing the assimilate supply. This happens when invertase, for example, is induced in leaves. Sink growth depends upon the amount of assimilates allocated to the specific sink. This can be stimulated by sink-
30 specific expression of the invertase. Since invertase activity negatively effects starch

synthesis, chemical control of invertase expression is clearly advantageous over its constitutive expression.

An increase sink supply is likely to result in larger tubers when invertase expression is induced. The early flowering phenotype is, however, believed to be explained by a
5 transient shortage of assimilate supply.

According to a second aspect of the present invention there is provided a method of controlling the flowering behaviour of a plant comprising transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to a controllable promoter
10 region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence(s) from said controllable promoter region by application of an external chemical inducer whereby the flowering behaviour of said transgenic plant is altered.

The method of controlling flowering behaviour may be used to speed up the growth
15 cycle of a plant such that more generations are produced.

The controllable promoter region in all aspects and embodiments of the present invention mentioned herein preferably comprises an inducible switch promoter system such as, for example, the alcA/alcR gene switch promoter system described in published International Patent Application No. WO 93/21334; the GST promoter as described in
20 published International Patent Application Nos. WO 90/08826 and WO 93/031294; and the ecdysone switch system as described in published International Patent Application No. WO 96/37609 the teachings of which are incorporated herein by reference. Such promoter systems are herein referred to as "switch promoters". Switch promoter systems are particularly suitable for use in the method of the present invention since they allow the
25 expression of DNA sequences to be switched on different parts of a transgenic plant at different times by means of sequential induction where the chemical inducer can be applied to the desired area of the plant at the desired stage of growth. For example, the switch chemical may be applied as a spray or vapour to all or part of the transgenic plant or as a root drench.

Examples of suitable switch chemicals are provided in the above references describing the switch promoter systems and are illustrated in the accompanying examples. The switch chemicals are desirably agriculturally acceptable chemicals.

Inducible promoter systems preferably include one or two component systems.

- 5 Systems comprising more than two components are, however, also included. The switch system may be driven by a constitutive promoter or, preferably, by a tissue or organ specific promoter whereby the target gene is only switched on in a target tissue or organ.

The alcA/alcR switch promoter system is particularly preferred for use in all aspects of the present invention mentioned herein.

- 10 The alcA/alcR inducible promoter system is a two component system involving DNA sequences coding for the alcA promoter and the alcR protein, the expression of which is placed under the control of desired promoters. The alcR protein activates the alcA promoter in the presence of inducer and any gene placed under the control of the alcA promoter will therefore be expressed only in the presence of inducer. The promoter controlling expression
15 of the alcR regulatory protein is preferably be a tissue or organ selective promoter, such as a leaf or tuber specific promoter, such that alcR is produced and alcA activated resulting in expression of the DNA sequence coding for the protein of interest only in selected parts of the plant such as for example the leaf, fruit, grain, endosperm or seed. When the method of the present invention is for use in cereal crops the expression of alcR is desirably controlled
20 by a seed specific promoter; for use in grain the expression of alcR is desirably controlled by promoters associated with genes involved in starch synthesis or with seed storage proteins and for use with forage crops the expression of alcR is desirably controlled by leaf specific promoters. Examples of tissue or organ selective promoters are well known in the art and include for example seed specific promoters such as the Ltp2 promoter (Kalla et al, Plant J 6
25 (6) 849-60, (1994)), the zmGBS, the zmZ27, the osAGP and the osGT1 promoters (Russell and Fromm, Transgenic Res 1997, 6 (2) 157-68), the CMd promoter (Grosset et al, Plant Mol Biol 1997 34(2) 331-338), the glycinin A2B1a promoter, (Itoh et al Mol Gen Genet 1994 243(3) 353-357), the oleosin promoter from Brassica napus (Keddie et al Plant Molecular Biology 19 443-453, (1992)), the MatP6 oleosin promoter from cotton (Hughes et al, Plant
30 Physiol (1993) 101 697-698), the oleosin promoter from Arabidopsis (Plant et al, Plant Mol. Biol. 25 193-205 (1994)), the zein promoter (Ottoboni et al Plant Mol Biol (1993) 21, 765-

778), and fruit and organ specific promoters such as the patatin promoter (Rocha-Sosa et al EMBO J 8 23-30 1989), the promoter family associated with ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit genes from tomato (Meier, Plant Physiol 107 (4) 1105-1118 (1995)), tomato rbcS3B and rbcS3C promoters (Carrasco Plant Mol Biol 21 (1) 1-15 (1993)), the leaf promoter STL1 (Eckes et al Mol. Gen Genet 205 14-22 (1986)) and the rolC promoters.

According to a further preferred embodiment of the present invention there is provided a method of increasing plant yield comprising transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to the alcA/alcR controllable promoter region wherein the promoter controlling expression of the alcR regulatory protein is a tissue or organ specific promoter and is optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence(s) from said controllable promoter region by application of an external chemical inducer whereby the yield of said transgenic plant is increased.

Examples of DNA sequences which may be used in the method of the present invention to increase plant yield and to control flowering behaviour include those DNA sequences coding for proteins involved in the transport, uptake and subsequent metabolism of sucrose e.g. phosphofructokinase, invertase and hexokinase; in sucrose biosynthesis e.g. sucrose synthase, sucrose phosphate synthase and fructose-1,6-bisphosphatase; in the transport of reserves during dormancy such as in phloem loading e.g. ATPase and sucrose and hexose transport proteins; in long distance phloem transport and in phloem unloading e.g. inorganic pyrophosphorylase (iPPase); in the utilisation of assimilates e.g. utilisation of sucrose-derived metabolites; in blocking starch synthesis (indirectly leading to increased sucrose levels); and invertase inhibitors.

The use of a controllable promoter region allows the production of the DNA sequence to be switched on in a controlled manner at the appropriate time in the growth cycle of the plant. We have unexpectedly found that the controlled expression of an invertase gene using the alcA/alcR switch promoter system leads to an increase in plant height, an increase in leaf size and to an increase of up to 10% in the fresh weight of a plant and accelerates the time at which the plants flower i.e the plants flower early.

According to a another preferred embodiment of the present invention there is therefore provided a method of increasing plant yield comprising transforming a plant with a DNA construct comprising a DNA sequence coding for an invertase operably linked to a controllable promoter region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence from said controllable promoter region by application of an external chemical inducer whereby the yield of said transgenic plant is increased.

According to yet another preferred embodiment of the present invention there is therefore provided a method of controlling the flowering behaviour of a plant comprising transforming a plant with a DNA construct comprising a DNA sequence coding for an invertase operably linked to a controllable promoter region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence from said controllable promoter region by application of an external chemical inducer whereby the flowering behaviour of said transgenic plant is altered.

The invertase may be derived from mammalian, bacterial, yeast, fungal or plant sources and may be different types such as acid invertase or neutral invertase. Invertase may be directed to different cellular locations such as the cell wall, the cytosol, the vacuole or apoplast by means of signal peptides (see Sonnewald *et al.* 1991 Plant J. 1:95-106).

According to a third aspect of the present invention there is provided a DNA construct comprising a DNA sequence coding for a protein involved in sucrose metabolism, uptake and/ or transport operably linked to a controllable promoter region.

The DNA constructs according to the present invention may also optionally contain a transcription terminator sequence and/ or a targeting sequence such that the invertase is targeted to a desired location within the plant. Examples of transcription terminators include the nopaline synthase transcription terminator and examples of suitable targeting sequences include for example signal sequences and vacuolar targeting sequences

In a preferred embodiment of this aspect of the present invention the DNA sequence codes for invertase and the controllable promoter region is an inducible promoter region comprising the *alcA/alcR* switch promoter system.

Plant cells may be transformed with recombinant DNA constructs according to a variety of known methods such as with *Agrobacterium* Ti plasmids, electroporation,

microinjection, microprojectile gun. The transformed cells may then be regenerated into whole plants in which the new nuclear material is stably incorporated into the genome. Some of the progeny of these primary transformants will inherit the recombinant DNA according to the present invention.

5 According to a fourth aspect of the present invention there is provided plant tissue transformed with a DNA construct comprising a DNA sequence coding for a protein involved in sucrose metabolism, uptake and or transport operably linked to a controllable promoter region and to the progeny of said plants.

10 Examples of suitable plants the yield of which may be increased and the flowering behaviour of which may be controlled according to the methods of the present invention and which may be transformed with DNA constructs according to the present invention include, for example, monocotyledonous and dicotyledonous plants such as field crops, cereals, fruit and vegetables such as : canola, sunflower, tobacco, sugarbeet, cotton, soya, maize, wheat, barley, rice, sorghum, tomatoes, mangoes, peaches, apples, pears, strawberries, bananas,
15 melons, potatoes, carrot, lettuce, cabbage and onion; trees such as eucalyptus and poplar trees and cut flowers and ornamentals.

20 The method of the present invention may be particularly useful for improving the uniformity of banana fillings in a hand of bananas where commonly the banana fingers at the top of the hand fill first and split while those at the bottom are not full enough. According to the method of the present invention the sink strength of the bananas may be altered such that fixed carbon from those at the top of the hand may be drawn into those bananas at the bottom leading to a more uniform hand size.

 The present invention is further illustrated only by way of example with reference to the following Examples and Figures in which :-

25 Figure 1 shows invertase activity in source leaves of transgenic Alc:cytosolic invertase and Alc: cell wall invertase tobacco plants following ethanol induction

 Figure 2 shows a graphical representation of quantum yield of wild type and transgenic tobacco plants at various time points after induction.

30 Figure 3 shows a histogram analysis of invertase activity in transgenic plants at two different concentrations of ethanol (i.e. for wild type, alc:INV27, alc:INV10, alc:INV28 and 35ScytINV

Figure 4 shows histogram analysis of a) invertase activity; b) fresh weight; c) height and d) % flowering plants in wild type and transgenic tobacco plants induced by ethanol (i.e. wild type, alc:INV27, alc:INV10 and alc:INV28)

Figure 5 shows a photograph of Alc invertase lines 27, 28, 10 and wild type

5 A) no ethanol B) with ethanol

Figure 6 shows transient expression of invertase increased flower number in Alc:cyINV but not Alc:cwINV tobacco plants

Figure 7 shows the alteration of flowering behaviour in both Alc:cyINV and Alc:cwINV on expression of invertase

10 Figure 8 shows transient expression of invertase leading to early flowering in tobacco plants

Figure 9 shows the draft strategy for cloning an Alc GUS construct with an L700 promoter

Figure 10 shows organ specific expression of L700::Alc:GUS tobacco plants after 48 hours induction

Figure 11 shows the draft strategy for cloning an Alc GUS construct with patatin B33

15 promoter

Figure 12 shows a histogram analysis of GUS activity in wild type and transgenic potato tubers 0 and 7 days after induction

Figure 13 shows a histogram analysis of the levels of GUS activity observed in tubers of potato transformed with a patatin:alc:GUS construct induced with ethanol

20 Figure 14 shows tissue specific and ethanol inducible GUS expression in transgenic tobacco plants: tuber-specific expression of the alcR protein

Figure 15 shows the plasmid construction of patatin B33::Alc:cwINV

Figure 16 shows invertase activity in patatin:Alc:cwINV potato tubers induced by ethanol

25 Figure 17 shows carbohydrate content in Pat::cwINV and Pat::Alc:cw INV potato growing tubers

Figure 18 shows increased potato tuber size resulting from early induction of apoplastic invertase expression

EXAMPLES

30 We have adapted the *alc* regulon of the ascomycete fungus *A. nidulans*, which has been well characterised (Pateman et al, Proc. Roy. Soc. London B 217, 243 (1983), Creaser

et al, Biochem J. 225 449 (1985), Lockington et al, Mol. Microbiol 1, 275 (1987), Felenbok et al, Gene 73, 385 (1988), Felenbok et al, J. Biotechnol. 17, 11 (1991), Kulmberg et al, J. Biol. Chem 267, 21146 (1992), Kulmberg et al, Mol. Microbiol. 7 847, (1993), and Fillinger and Felenbok, Mol. Microbiol 20 475 (1996)). From classical genetics, it has been assumed
5 that this is a self-contained genetic system that controls the cellular response to ethanol and other related chemicals. In *A.nidulans*, the *alcA* and *aldA* genes encode alcohol dehydrogenase I and aldehyde dehydrogenase, respectively (Pateman et al, Proc. Roy. Soc. London B 217, 243 (1983), Creaser et al, Biochem J. 225 449 (1985), Lockington et al, Mol. Microbiol 1, 275 (1987), Felenbok et al, Gene 73, 385 (1988), Felenbok et al, J. Biotechnol.
10 17, 11 (1991), Kulmberg et al, J. Biol. Chem 267, 21146 (1992), Kulmberg et al, Mol. Microbiol. 7 847, (1993), and Fillinger and Felenbok, Mol. Microbiol 20 475 (1996)). Both of these genes are regulated by the pathway-specific AlcR transcription factor (Pateman et al, Proc. Roy. Soc. London B 217, 243 (1983), Creaser et al, Biochem J. 225 449 (1985), Lockington et al, Mol. Microbiol 1, 275 (1987), Felenbok et al, Gene 73, 385 (1988),
15 Felenbok et al, J. Biotechnol. 17, 11 (1991), Kulmberg et al, J. Biol. Chem 267, 21146 (1992), Kulmberg et al, Mol. Microbiol. 7 847, (1993), and Fillinger and Felenbok, Mol. Microbiol 20 475 (1996)). The AlcR protein binds to specific sites within the *alcA* promoter region and, as we demonstrate here, must respond directly to the inducer molecule (Pateman et al, Proc. Roy. Soc. London B 217, 243 (1983), Creaser et al, Biochem J. 225 449 (1985),
20 Lockington et al, Mol. Microbiol 1, 275 (1987), Felenbok et al, Gene 73, 385 (1988), Felenbok et al, J. Biotechnol. 17, 11 (1991), Kulmberg et al, J. Biol. Chem 267, 21146 (1992), Kulmberg et al, Mol. Microbiol. 7 847, (1993), and Fillinger and Felenbok, Mol. Microbiol 20 475 (1996)).

The *alc* regulon was considered suitable for a plant gene expression cassette for a
25 number of reasons. First, the minimal regulon would include only the *alcR* gene and the *alcA* promoter. Second, the evolutionary divergence between *A. nidulans* and higher plants would make it unlikely that any plant homologues of the AlcR protein would activate the promoter: AlcR contains a zinc binuclear cluster like Gal4 (Pateman et al, Proc. Roy. Soc. London B 217, 243 (1983), Creaser et al, Biochem J. 225 449 (1985), Lockington et al, Mol. Microbiol
30 1, 275 (1987), Felenbok et al, Gene 73, 385 (1988), Felenbok et al, J. Biotechnol. 17, 11 (1991), Kulmberg et al, J. Biol. Chem 267, 21146 (1992), Kulmberg et al, Mol. Microbiol. 7

847, (1993), and Fillinger and Felenbok, Mol. Microbiol 20 475 (1996)) which has only been found in fungi so far. In addition, no other plant transcription factors were likely to cause interference at the *alcA* promoter. Third, the chemical inducers are relatively simple organic molecules with low phytotoxicity. Fourth, under normal growth conditions, the levels of
5 natural inducers in the plant would be extremely low.

To test the efficacy of the system, plant expression cassettes were constructed. Construction of the *alc* derived gene constructs. p35S: *alcR* (A) utilised the 35S promoter from the Cauliflower Mosaic Virus to express AlcR protein from a cDNA. A partial *alcR* cDNA (provided by Felenbok) was excised from its Bluescript vector (Stratagene) by partial
10 digestion with *Bam*HI, ligated to *Bam*HI digested pJR1 (Smith et al, Nature 334, 724 (1988)), a pUC derived vector containing the CaMV 35S promoter and the *nos* terminator, and transformed into *E. coli* XL-1 Blue (W.O. Bullock et al. BioTechniques 5, 376 (1987); J. Sambrook et al., Molecular cloning: A laboratory manual, edn. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, 1989). The *alcA* reporter cassette, *palcA*:
15 CAT (B), was constructed by digestion of pCaMVCN with *Hind*III and *Bam*HI to remove the promoter. (pCaMVCN is a plant expression vector available from Pharmacia. It is a pUC - derived vector in which the CaMV 35S promoter expresses the bacterial CAT gene. The terminator is from the *nos* gene of *A. tumefaciens*.) Since the TATA boxes of the *AlcA* and 35S promoters were identical (5'TCTATATAA3'), recombinant PCR was used to
20 amplify and fuse both fragments through this site (Higuchi in PCR Protocols, M.A. Innis et al, eds (Academic Press, San Diego (1990) p177-183). The *alcA* PCR product extended from the TATA box for 246 bp upstream, and included ALCR binding sites (Pateman et al, Proc. Roy. Soc. London B 217, 243 (1983), Creaser et al, Biochem J. 225 449 (1985), Lockington et al, Mol. Microbiol 1, 275 (1987), Felenbok et al, Gene 73, 385 (1988),
25 Felenbok et al, J. Biotechnol. 17, 11 (1991), Kulmberg et al, J. Biol. Chem 267, 21146 (1992), Kulmberg et al, Mol. Microbiol. 7 847, (1993), and Fillinger and Felenbok, Mol. Microbiol 20 475 (1996)). The 35S PCR product included the common TATA box sequence and extended downstream to span a convenient *Bam*HI site for subsequent cloning; the minimal 35S promoter is known not to be expressed in plants (It has been shown that a
30 minimal 35S promoter containing only those sequences between positions -46 and +5 lacks the ability to initiate transcription (Odell et al Nature 346, 390 (1985); Hilson et al, Plant Cell

2 651 (1990) Schena et al Proc. Natl. Acad Sci USA 88, 10421 (1991). It is reasonable to expect that fusion of *palcA* through the TATA sequence (positions -31 to +1) would also be inactive.) The resultant product was digested with *Hind*III and *Bam*HI, and ligated into the promoterless pCaMVCN. After transformation into *E.coli*, colonies were screened to select a
5 plasmid which contained the appropriate *palcA*:35S fusion promoter, and the *Hind*III and *Bam*HI fragment was sequenced to ensure that there were no PCR errors. The *palcA*::*Inv* construct was obtained by deletion of the GUS reporter gene from plasmid *palcA*:GUS and insertion of the truncated yeast *suc2* gene isolated from plasmid *rolC-suc2* as a *Bam*HI fragment (Lerchl et al (1995) Plant Cell 7, 259 (1995). For plant transformation, the
10 p35S:*alcR* cassette was cloned into a Bin19-derived vector (Deblacre et al, Nucleic Acids Res. 13, 4777 (1985), together with either the *palcA*:CAT or the *palcA*::*Inv* construct, transformed into *A. tumefaciens* (Holsters et al. Mol. Gen. Genet. 163, 181 (1978); Vervliet et al. J. Gen. Virol 26, 33 (1975)). Tobacco transforming using *Agrobacterium*-mediated gene transfer was carried out as described previously (Rosahl et al EMBO J. 6:1159 (1987)
15 and Komari et al Plant Science 60 223 (1989)).

The bacterial chloramphenicol acetyltransferase gene (CAT) was used as a reporter gene so that levels of expressed protein could be determined using ELISA. When transformed into *A. nidulans* (Ballance and Turner Gene 36, 321 (1983); Campbell *et al*). Curr. Genet. (1989), the *palcA*:CAT construct showed inducible CAT activity, and
20 p35S:*alcR* restored the wild-type phenotype to an *alcR* mutant (data not shown). Transient assays (Callis et al Genes and Develop 1 1183 (1987)) in maize protoplasts revealed that the AlcR protein could stimulate the transcription from the *alcA* promoter in plant cells and that expression was at least partially regulated by ethanol (data not shown).

After *Agrobacterium tumefaciens* mediated transformation a transgenic tobacco plant
25 carrying the p35S:*alcR* and *palcA*:CAT cassettes was selected and tested by PCR for the presence of both cassettes (data not shown). This plant was selfed, and the seedling progeny assayed for both the selectable marker and CAT expression. The construct segregated among the progeny in a Mendelian ratio (1 non-transgenic: 2 hemizygous: 1 homozygous) consistent with a single copy of the cassettes in the parent plant (data not shown).

30 A selected seedling was grown to maturity to produce a homozygous line. Seedlings of this plant were tested for CAT protein in comparison to seedlings of a similar plant

transformed with a construct which expressed CAT from the constitutive high activity CaMV 35S promoter (Table 1). The homozygous *palc*:CAT seedlings had barely detectable CAT protein in the absence of induction, but had 39% of the CAT activity of the untreated p35S:CAT seedlings and 55% relative to the ethanol-treated p35S:CAT seedlings. Thus, ethanol treatment of p35S:CAT seedlings resulted in a reduction [29%] in CAT protein levels relative to the untreated control.

While the induced levels of expression were lower than that from the 35S promoter, the negligible basal activity indicated its suitability for the manipulation of carbon metabolism. A range of inducible invertase vectors were made by replacing the CAT reporter gene with the truncated yeast *suc2* GENE 100, a cytosolic yeast derived invertase. In this regard, the Bam HI fragment isolated from plasmid RolC::Suc2 (Lerchl *et al.*, 1995, Plant Cell 7, 259-270) was used to replace the reporter gene. The fragment contained nucleotides 848 to 2393 of the yeast *suc2* gene (Accession number Y01311) and encoded an invertase protein without a signal peptide. Invertases from other sources and of different type, such as acid invertase, or other targeted invertases were also made using transit peptide invertase combinations described in Sonnewald *et al.*, 1992, Plant J. 1:95-106 which may be expressed in the cell wall or subcellular locations such as the vacuole or the apoplast (details of cloning are described in Caddick *et al.* Nature Biotech, Vol 16, Feb 1998 page 177 and in Lerchl J. *et al.* 1995, Plant Cell 7: 250-270 and Sonnewald *et al.*). Transgenic tobacco plants carrying *palcA*:cyInv were isolated (Tobacco (*Nicotiana tabacum* cv. Samsun NN) transformation using *Agrobacterium*-mediated gene transfer was carried out as described by Rosahl *et al.* EMBO J. 13, 1 (1987)). After screening about 100 independent kanamycin resistant regenerants for ethanol inducible invertase activity, 23 invertase expressing plants could be identified. Of 23 plants exhibiting inducible invertase activity, three lines [10, 27 and 28] were selected for more detailed analysis. To this end, plants were multiplied in tissue culture and 50 plants of each line were transferred into the greenhouse. After 21 days of growth in 2L pots, initial induction was carried out via root drenching with 100ml of a 1% ethanol solution (v/v). To accelerate the ethanol response, induction was repeated at 48 and 72 hours after the initial root drench. To assay invertase activity, samples were taken at 0, 1, 6, 24, 48, 72 and 96 hours after the initial induction (see Figure 1). Elevated invertase activity was measurable in all three transgenic lines already 6 hours after the first addition of

ethanol. Invertase activity increased steadily reaching a plateau 96 hours after the initial root drenching in two lines [10 and 27], while in the third [28], it was still increasing.

Phenotypic modification started 72 hours after ethanol induction and was strongest after 96 hours. The final phenotype was identical to the previously published results using the 35S CaMV promoter to drive the expression of cytosolic yeast invertase (Sonnewald et al
5 the 35S CaMV promoter to drive the expression of cytosolic yeast invertase (Sonnewald et al Plant J. 1, 95 (1991)). Development of this phenotype followed maximal invertase activity and was most severe in transformant 28. Photosynthesis fluorescence measurements were used to monitor changes of quantum yield (Schreiber et al, in Ecophysiology of Photosynthesis Vol. 100, Schuize and Caldwell, Eds (Springer Verlag, Berlin, 1994), pp 49-
10 70.) of all three transformant lines *in vivo* throughout the induction experiment. During the course of ethanol treatment, quantum yield did not change markedly in the youngest leaves (leaves A 8% of maximal leaf area). However, coinciding with the developing visual phenotype quantum yield decreased significantly ($p > 0.05$) in leaves B (15% of max) and C (45% of max) of plants from line 10 and 28 starting 72 hours after the initial induction and
15 developing further until the final time point at 96h.

Figure 2 shows evidence for a reduced rate of photosynthesis following the increase of invertase activity in transgenic tobacco plants as determined by quantum yield measurements. Fluorescence measurements were used to monitor changes in photosynthetic parameters during induction of invertase activity using the PAM-2000 instrument (Walz,
20 Effeltrich, Germany). Quantum use efficiency (quantum yield) of photosystem II (PSII) was measured by applying a saturating light beam on light adapted leaves of wildtype and transformed plants (palc:Inv). Before each measurement, it was verified that the saturating pulse had reached a plateau to allow an accurate determination of F_m' . The intensities of the measuring and saturating light beam were adjusted to reach a FO' value close to 0.4.
25 Measurements were conducted on different leaves having reached 8% (A), 15% (B), or 45% (C) of maximal leaf area of five plants of each genotype at the indicated time points.

Quantum yield of three succeeding light adapted leaves (leaf A-C) starting from the top of the plant was measured using a PAM-2000 instrument at the time points indicated. Values given are the means \pm SE ($n=5$). For plants of line 28, quantum yield was reduced by
30 23% ($p < 0.05$) and 27% ($p < 0.05$) and for plants of line 10 only by 6% and 17% ($p < 0.05$),

respectively. Due to heterogeneity of the developing phenotype between individual plants from each genotype, standard errors were higher in the affected leaves (B and C).

The table below shows CAT activity levels in transgenic tobacco. Individual seedlings from a homozygous transgenic tobacco line carrying the CAT gene expressed from the *alcA* promoter were compared with those from a similar line transformed with p35S:CAT. Plants were grown on liquid media until 4 weeks old, and showed four true leaves (seed progeny of tobacco plants were grown by sowing seeds directly onto a 2cm layer of sterile alcatene beads (5mm diameter) floating on a sterile solution of 0.5% (w/v) Miracle Gro in 500ml beakers. The beakers were covered with a perforated plastic bag and incubated at 25°C under high intensity lights in a growth room). Induction was achieved by the addition of 0.1% ethanol to the growth medium for 120 h. The induction medium was replaced at 58 h to maintain ethanol concentrations. One leaf was taken prior to induction, and one leaf after induction. CAT ELISA (Boehringer Mannheim) was performed on crude cell extracts; total protein was determined as described previously. (Bradford, Anal Biochem 72: 243 (1976). All values are ng CAT protein per mg total protein, and represent the mean of nine individual replicates \pm one standard deviation.

<u>Line</u>	<u>Untreated</u>	<u>Ethanol-induced</u>
palcA:CAT	0.36+0.43	30.37+3.91
p35S:CAT	78.08+30.44	55.46+10.85

It can be seen from Figure 3 that the invertase activity in transgenic plants is dose-dependent and that the activity at 5% ethanol is significantly greater than it is at 1% ethanol. It is, therefore, possible to regulate invertase in a dose dependent manner using the Alc switch.

In order to see the impact of inducible cytosolic invertase expression on plant growth and flowering time, tobacco plants were vegetatively propagated in tissue culture (Figure 4). Subsequently 50 plantlets each genotype were transferred into the greenhouse. In Figure 4, wt denotes a wild type transgenic control and lines 10, 27 and 28 represent 50 cloning propagated independent lines containing 35S:alc A:suc 2. Three weeks after transfer, plants were induced with 100ml 1% (v/v) via root drenching. Induction was repeated three times

(0, 48 and 72 hours). In particular, Figure 4a) shows cytosolic neutral invertase activity (suc 2) measured 96 hours after initial induction, Figure 4b) shows fresh weight of the above ground biomass 45 days after transfer and Figure 4c) shows plant height 45 days after transfer. Figure 4d) shows the percentage of plants which were flowering when scored 45
5 days after transfer.

To show the impact of ethanol inducible cytosolic invertase on plant height and flowering, plants were propagated in tissue culture and transferred into the greenhouse (Figure 5). Three weeks after transfer one half of the plants were induced with ethanol as described for Figure 4. The second half of the plants was transferred into a second
10 greenhouse without any ethanol treatment. The upper panel (A) shows four tobacco plants 7 weeks after transfer from tissue culture without ethanol induction. The lower panel (B) shows the same genotypes 4 weeks after initial ethanol induction. From left to right the following genotypes are shown: 1, line 27; 2, line 10; 3, line 28; line 4, untransformed control. The early flower phenotype was consistently found in all experiments.

15 In order to show that inducible invertase expression leads to an increased flower number per plant, 25 plants of each genotype (wt, cyt inv10, cy inv 27, cyt inv 28, cw inv 19, cw inv 28 and cw inv 45) were propagated in tissue culture and transferred into the greenhouse (Figure 6). Three weeks after transfer plants were induced as described above. The total number of flowers produced each plant was determined at the end of the growing
20 period.

As can be seen from Figure 7, transgenic plants expressing inducible invertase have accelerated flower induction. 25 plants of each genotype were propagated in tissue culture and transferred into the greenhouse. Three weeks after transfer plants were induced as described above. Subsequently, flower formation was followed throughout the growing
25 period. Plants were classified as flowering when the first flower bud was open. Values are given in [%] flowering plants per total number of plants (n=25).

As can be seen from Figure 8, the early flower phenotype is reproducible at different growing seasons by means of transient expression of invertase. In this regard, 50 (spring) and 25 (summer and autumn) plants of each genotype were used for the individual
30 experiments, respectively. After propagation, plants were transferred into the greenhouse and induction started three weeks after transfer as described above. At the indicated time

after transfer (dpt, days after transfer) plants with open flower buds were counted. Values are given in [%] flowering plants per total number of plants.

Preparation of Plasmid pSTLS1:AlcR:AlcA:GUS (SC08)

- 5 To obtain plasmid SC08, the EcoRI/HindIII fragment of plasmid AlcR/A GUS containing the AlcR coding region and the NOS terminator was subcloned into pBluescript SK-yielding plasmid pAlcR. Subsequently, plasmid pAlcR was digested with EcoRI, blunt ended with DNA polymerase (Klenow fragment), further restricted with HindIII and ligated into plasmid pBINSTSL1 after BamHI digestion, Klenow treatment and Hind III digestion yielding
- 10 plasmid pBIN:STSL1:AlcR. Plasmid pBINSTSL1 consists of the STSL1 promoter corresponding to nucleotide +1 to +1585 of the published sequence of the STSL1 gene from potato (Eckes et al (1986) Mol. Gen. Genet. 205 14-22) and the OCS (octopine synthase) terminator. The final construct SC08 was obtained by ligating the HindIII fragment from plasmid AlcR/A GUS containing the AlcA promoter, the GUS coding region and the NOS
- 15 terminator into the HindIII digested vector pBIN:STSL1:AlcR. The strategy for cloning an Alc GUS construct with the L700 promoter is shown in Figure 9.

- The tissue specific and ethanol inducible GUS expression in transgenic tobacco plants i.e. leaf/stem-specific control of the alcR protein is shown in Figure 10. Transgenic tobacco plants expressing the GUS reporter gene under control of the ethanol inducible
- 20 system were propagated in tissue culture and transferred into the greenhouse. Three weeks after transfer plants were induced via root drenching using 100ml 1% ethanol. 48 hours after induction tissue samples were harvested and GUS activity determined in protein extracts:- sink leaves, < 3cm; source leaves. 35S::Alc:GUS, root and stem expression of the alcR protein under control of the 35S CaMV promoter was used as a constitutive control.
- 25 L700:: Alc:GUS expression of the alcR protein under control of the leaf/stem-specific ST-LS1 promoter from potato in 4 independent transgenic lines (6, 9, 27 and 74) were also used.

Preparation of Plasmid B33:AlcR:AlcA:GUS (SC09)

- To obtain plasmid SC09, the EcoRI/HindIII fragment of plasmid AlcR/A GUS containing
- 30 the AlcR coding region and the NOS terminator was subcloned into pBluescriptSK- yielding plasmid pAlcR. Subsequently, plasmid pAlcR was digested with EcoRI, blunt ended with

DNA polymerase (Klenow fragment), further restricted with HindIII and ligated into the SmaI/HindIII digested plasmid pBIN:B33AlcR. Plasmid pBINB33 consists of the patatin class I promoter, corresponding to nucleotide -1512 to +14 of the patatin gene B33 (Rocha-Sosa et al. (1989) EBO J. 8 23-29) and the OCS terminator. The final construct SC09 was
5 obtained by ligating the HindIII fragment from plasmid AlcR/A GUS containing the AlcA promoter, the GUS coding region and the NOS terminator into the HindIII digested vector pBIN:B33AlcR. A strategy for cloning an Alc GUS construct with a patatin B33 promoter is shown in Figure 11.

10 Alc R patatin alc A GUS vector

pSC09 (B33-alc GUS in Bin 19) was transformed directly into *Agrobacterium tumefaciens* strain C58C1:pGV2260 using the protocol described by Hofgen and Willmitzer (1988). Potato (var Solara) transformation using *Agrobacterium*-mediated gene transfer was performed as described by Roscha-Sosa et al (1989). Transgenic plants were duplicated in
15 tissue culture and one set transferred to the glasshouse following root formation. Plants were grown to maturity and tubers harvested. For each independent transformant tuber, samples were taken for GUS analysis in the absence of ethanol treatment. Additional tubers were transferred to perspex boxes containing a pot of 1% ethanol. Following 7 days of ethanol vapour treatment, tubers were harvested and assayed for GUS activity. Figure 13
20 demonstrates that high levels of transgene expression is observed in the tubers following ethanol treatment.

Transgenic potato plants expressing the GUS reporter gene under control of the ethanol inducible system were propagated in tissue culture and transferred into the greenhouse. Two months after transfer, plants were induced via root drenching using 100ml
25 1% ethanol. 48 hours after induction, tissue samples were harvested and GUS activity determined in protein extracts:- leaves, stems and tubers > 5g. Pat:GUS expression of the GUS reporter gene under control of the class I patatin promoter B33 acted as a control. Pat::Alc:GUS expression of the alcR protein under control of the tuber-specific B33 promoter from potato was also used. The activity is given in pmol MU/mg/min. Figure 14
30 shows tissue specific and ethanol inducible GUS expression in tobacco plants by tuber-specific expression of the alcR protein.

The EcoRI and HindIII fragment containing transactivator AlcR gene and NOS terminator was from plasmid 35S:AlcR-AlcA:GUS (35S-Alc:GUS, Plant Journal 1998, 16 (1) 127-132) and subsequently subcloned into Bluescript SK- (STRATAGENE) (AlcR in SK-). The plasmid AlcR in SK- was digested with EcoRI and filling-in with Klenow
5 fragment to make it blunt and further cut with HindIII. This EcoRI(-) - HindIII fragment was cloned into binary vector Bin-B33 cut by Small and HindIII resulting in the plasmid patatin B33:AlcR in pBIN19. The PCR product of yeast invertase with protein inhibitor II signal peptide (SP) sequence (von Schaewen *et al.* (1990) EMBO J. 9. 3033-3044) was cloned into pGEM-T vector by primer K83 and K84 which containing the Small site. The Small
10 fragment was subcloned into pUC-AlcA plasmid via BamHI site which was blunted by T4 DNA polymerase. The orientation was checked by combining Asp718 and EcoRI, and also with XbaI. The correct orientation plasmid was subsequently subcloned into patatin B33:AlcR in pBIN19 resulting the final plasmid patatin B33:AlcR-AlcA:cwINV in BIN19 (patatin B33::Alc:cwINV). The construction of the plasmid is shown in Figure 15.

15 Transgenic potato plants were propagated in tissue culture and subsequently transferred into the greenhouse. After tuber setting, plants were induced once with 100ml-1% ethanol (root drenching) and invertase activity determined in tubers. The left panel shows invertase activity prior to (0) and after (1) ethanol induction visualised after SDS-PAGE. An untransformed wild type was used as a control and compared against
20 independent transgenic lines 2, 3, 4, 5, 7 and 13 (see Figure 16).

Figure 17 shows the carbohydrate content of potato tubers two months after initial induction. Pat::cwINV, transgenic plants expressing yeast invertase were placed under control of the tuber-specific B33 patatin promoter (independent transgenic lines 3, 33 and 41). SC12 (Pat:: Alc:cwINV) transgenic plants (independent transgenic lines 2, 5, 7 and 13)
25 are ethanol inducible. Tuber-specific expression of cell wall invertase is caused via tuber-specific expression of the alcR protein mediated by the B33 promoter.

Transgenic potato plants were propagated in tissue culture and transferred into the greenhouse. Ethanol induction occurred at three different development stages. 1st induction, 25 days after transfer, 2nd induction 32 days after transfer and a 3rd induction 39 days after
30 transfer. 10 plants from each genotype were used for each induction experiments. Initial induction occurred via root drenching. Due to the induction procedure plants induced after

25 days were vapour induced 32 and 39 days after transfer. Plants induced 32 days after transfer were induced a second time, whereas, plants induced 39 days after transfer were induced only once (see Figure 18).

Other modifications of the present invention will be apparent to those skilled in the
5 art without departing from the scope of the invention.

CLAIMS

1. A method of increasing the yield of a plant comprising transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to a controllable promoter region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence(s) from said controllable promoter region by application of an external chemical inducer whereby the yield of said transgenic plant is increased.
2. A method of increasing the yield of a plant according to claim 1 by selectively increasing the importation of fixed carbon into photosynthetically inactive sink tissue comprising transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to a controllable promoter region and optionally operably linked to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence(s) from said controllable promoter region by application of an external chemical inducer whereby the transportation of fixed carbon from photosynthetically active source tissue to photosynthetically inactive tissue of said transgenic plant is selectively increased.
3. A method of controlling the flowering behaviour of a plant comprising transforming a plant with a DNA construct comprising one or more DNA sequence(s) coding for a protein involved in sucrose sensing, transport, metabolism and/or uptake operably linked to a controllable promoter region and optionally to a transcription terminator and controlling the level, time and spatial location of expression of said DNA sequence(s) from said controllable promoter region by application of an external chemical inducer whereby the flowering behaviour of said transgenic plant is altered.
4. A method according to any one of the preceding claims wherein said DNA sequence(s) include those DNA sequences coding for proteins involved in the

transport, uptake and subsequent metabolism of sucrose, such as in sucrose biosynthesis; in the transport of reserves during dormancy such as in phloem loading; in long distance phloem transport and in phloem unloading; and in the utilisation of assimilates such as sucrose-derived metabolites.

5

5. A method according to claim 4 wherein said DNA sequence(s) include those coding for sucrose synthase, phosphofructokinase, invertase, hexokinase, inorganic pyrophosphorylase or ATPase.

10

6. A method according to claim 5 wherein the DNA sequence codes for an invertase.

7. A method according to any one of the preceding claims wherein said controllable promoter region comprises a chemically inducible promoter system.

15

8. A method according to claim 7 wherein the chemically inducible promoter is under the control of a tissue or organ selection promoter.

9. A method according to claim 7 or claim 8 wherein said chemically inducible promoter system comprises the *alcA/alcR* promoter system.

20

10. A method according to claim 9 wherein expression of the *alcR* regulatory protein is under the control of a tissue or organ selective promoter.

25

11. A DNA construct comprising a DNA sequence(s) coding for a protein involved in sucrose metabolism, uptake and or transport operably linked to a controllable promoter region.

12. A DNA construct according to claim 11 wherein said controllable promoter region comprises a chemically inducible promoter system.

30

13. A DNA construct according to claim 12 wherein said chemically inducible promoter system is the alcA/alcR switch promoter system.
14. A DNA construct according to claim 13 wherein the alcR regulatory protein is under
5 the control of a tissue or organ selective promoter.
15. A DNA construct according to any one of claims 11 to 14 wherein said DNA sequence(s) comprise a DNA sequence coding for an invertase.
- 10 16. Plant tissue transformed with a DNA construct according to any one of claims 11 to 15 and the progeny of said plants.
17. A method or a DNA construct substantially as hereinbefore described with reference to any one of the Figures .




Fig.2.





Fig.3.



4/16

Fig. 4.




Fig.5.


A



B





Fig.6.



7/16

Fig.7.



8/16

Fig.8.





Fig.9.



10/16

Fig.10.





Fig.11.



12/16

Fig.12.




Fig.13.




Figure 1 is a bar chart showing GUS activity (pmol MU/mg/min) for two constructs, Pat::GUS and Pat::AlcGUS, across five tissues (Leaves, Stems, Stolons, Tubers, Roots) under two conditions: Uninduced (U) and Ethanol treated (I). The Y-axis ranges from 0 to 45,000 pmol MU/mg/min. The X-axis lists tissues and conditions. Error bars represent standard deviation.

Construct	Tissue	Condition	GUS activity (pmol MU/mg/min)
Pat::GUS	Leaves	U	~100
		I	~100
	Stems	U	~100
		I	~100
	Stolons	U	~100
		I	~100
	Tubers	U	~32,000
		I	~32,000
	Roots	U	~100
		I	~100
Pat::AlcGUS	Leaves	U	~100
		I	~100
	Stems	U	~100
		I	~100
	Stolons	U	~100
		I	~100
	Tubers	U	~100
		I	~100
	Roots	U	~100
		I	~100

Fig. 15.

Restriction map of the pNOS-35S plasmid. The map shows a circular DNA molecule with several restriction sites and fragments. The fragments are labeled: 35S, A16R, NOS, A16A, SP, INV, and NOS. The restriction sites are: EcoRI, BamHI, HindIII, SmaI, and XbaI. The map is divided into two main sections by a vertical line. The left section contains the 35S, A16R, and NOS fragments. The right section contains the A16A, SP, INV, and NOS fragments. The restriction sites are indicated by vertical lines with labels above or below them.

Fig.16.




Fig.17.





Fig.18.



INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03687

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 C12N15/82 C12N9/26 A01H5/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 C12N A01H Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 07221 A (RIESMEIER JOERG ;TRETWEY RICHARD (DE); WILLMITZER LOTHAR (DE)) 27 February 1997	1-8, 11, 12, 15-17
Y	see the whole document	9, 10, 13, 14
Y	--- WO 93 21334 A (CADDICK MARK XAVIER ;ZENECA LTD (GB); GREENLAND ANDREW JAMES (GB);) 28 October 1993 cited in the application see the claims see abstract; figures 1-3; examples 1-5 see page 3 - page 16	9, 10, 13, 14
X	--- EP 0 442 592 A (INST GENBIOLOGISCHE FORSCHUNG) 21 August 1991 see the whole document --- -/-	1-8, 11, 12, 15-17
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
<p>* Special categories of cited documents :</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
Date of the actual completion of the international search 18 March 1999		Date of mailing of the international search report 25/03/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Oderwald, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 98/03687

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 438 904 A (CAMBRIDGE ADVANCED TECH) 31 July 1991 see the whole document -----	1-6, 11, 15-17
P, X	CADDICK M X ET AL: "AN ETHANOL INDUCIBLE GENE SWITCH FOR PLANTS USED TO MANIPULATE CARBON METABOLISM" NATURE BIOTECHNOLOGY, vol. 16, February 1998, pages 177-180, XP000764088 see the whole document -----	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 98/03687

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9707221 A	27-02-1997	DE 19529696 A	13-02-1997
		AU 6820496 A	12-03-1997
		CA 2229061 A	27-02-1997
		CN 1196090 A	14-10-1998
		EP 0846180 A	10-06-1998
WO 9321334 A	28-10-1993	AU 3901993 A	18-11-1993
		EP 0637339 A	08-02-1995
EP 0442592 A	21-08-1991	DE 4004800 A	14-08-1991
		AU 650639 B	30-06-1994
		AU 7089891 A	15-08-1991
		CA 2036103 A	14-08-1991
		JP 5049482 A	02-03-1993
		US 5436394 A	25-07-1995
EP 0438904 A	31-07-1991	AU 6836590 A	04-07-1991
		JP 4341126 A	27-11-1992
		KR 123183 B	20-11-1997
		US 5387756 A	07-02-1995